

# *The Risks of Diving While Pregnant*

## Reviewing the Research



David Haas Photo

### **Should a pregnant woman scuba dive?**

Whether expectant women should dive is a question that affects not only female divers but also their partners, dive buddies and dive professionals. Most divers can recall from their open water training that women are encouraged to stop diving during pregnancy, but few classes go into further detail.

What are the risks of diving while pregnant? What is it about scuba diving that is dangerous for a developing foetus? The published literature provides a foundation for the discussion.

**BY HEATHER E. HELD, BS AND  
DR. NEAL W. POLLOCK, PHD**

As with all research, there are limitations on how much the available studies can tell us. For ethical reasons, experiments with pregnant women are very limited. Most studies conducted with humans are surveys, and surveys have weaknesses, most importantly that they are not as easily controlled as laboratory research and that they can easily be biased.

A survey of female divers who had recently given birth included 69 women who had not dived during their pregnancies and 109 women who had. The nondiving women reported no birth defects, while the diving women reported an incidence of 5.5 percent.<sup>7</sup> To provide perspective, the survey author stated that the latter rate was within the normal range for the national population. The small sample size and the likelihood of selection bias in those responding to the survey make the results even more difficult to interpret.

While surveys can establish correlations, they cannot confirm causal relationships. In this case, they cannot confirm that diving caused a defect. To obtain such data, scientists rely on more highly controlled animal studies.

### 'Diving' in chambers

Hyperbaric chambers, which can simulate the increased pressure of diving, have been used to test different species of animals. Those results must then be translated to the human experience.

Many complex processes occur during pregnancy, and insults (disruptions of normal events) can lead to varied complications. Most diving-related studies have addressed the first and third trimesters of pregnancy. First trimester research has concentrated on the teratogenic, or birth-defect-causing, effects of hyperbaric oxygen (HBO). Third trimester research has examined the effects of decompression sickness (DCS) on the foetus and how diving

**TABLE 1: Depth (approx) at oxygen partial pressure limits for humans**

Breathing Gas	PO <sub>2</sub> Limit	
	1.4 ATA Depth (msw)	1.6 ATA Depth (msw)
100% O <sub>2</sub>	4	6
36% Nitrox	29	35
32% Nitrox	34	40
Air (21% O <sub>2</sub> )	57	66

and the foetal circulatory system interact.

A range of developmental abnormalities have been associated with hyperbaric exposure. These include low birth weights among the offspring of diving mothers<sup>14,15,26</sup>; foetal abortion<sup>28</sup>; bubbles in the amniotic fluid<sup>13,25</sup>; premature delivery<sup>14</sup>; abnormal skull development<sup>1,15,16</sup>; malformed limbs<sup>11,15,16</sup>; abnormal development of the heart<sup>16,20</sup>; changes in the foetal circulation<sup>2</sup>; limb weakness associated with decompression illness<sup>21</sup>; and blindness<sup>14</sup>.

We expose ourselves to hyperbaric oxygen - that is, oxygen concentrated by pressure - during almost all dives. A safe limit for the partial pressure of oxygen (PO<sub>2</sub>) is frequently accepted as 1.4 to 1.6 atmospheres of absolute pressure (ATA)<sup>19</sup>. Table 1 shows the depth (in msw) where these PO<sub>2</sub> levels are achieved with different breathing gas mixtures.

Rodents, which have large litters and relatively short gestational periods<sup>12</sup>, have been used to study the effects of HBO on developing foetuses. Female hamsters experiencing untreated DCI had offspring with severe limb and skull abnormalities.<sup>15,16</sup> Pregnant hamsters experiencing HBO-treated decompression illness also bore offspring with defects, though with less frequency than the untreated group<sup>15</sup>. Neither study reported noticeable differences in anatomical development between offspring from the nondiving control group and the group that dived without developing signs of DCI<sup>15,16</sup>.

Foetal rat hearts have proven sensitive to multihour HBO exposure (3.0 ATA for eight hours), albeit of a magnitude in excess of what humans could tolerate. In almost half the cases, the septum, which divides the right and left sides of the heart, failed to form properly<sup>20</sup>. Major blood vessels were positioned incorrectly just as often, compromising normal circulatory patterns<sup>20</sup>.

Another study of HBO-exposed rats found no significant differences between offspring from mothers that had dived and offspring from mothers that had not dived<sup>6</sup>. The PO<sub>2</sub> in this study (1.3 ATA for 70 minutes) was significantly less than that used in the previous study. The treatment difference may explain the dissimilar results.

Table 2 shows a summary of the timeline for human foetal development. It appears that hyperbaric exposure can alter the signals foetal tissues rely on to correctly orchestrate developmental processes. The nature of the abnormality is influenced by the timing of the insult. It is important to note, however, that exposure will not affect development in all instances.

### Decompression stress

The relative risk of decompression stress on mother and foetus is another question for consideration. Given sufficient decompression stress, blood returning to the heart from the body may contain venous gas emboli (VGE or bubbles)<sup>23</sup>. Sheep have been studied frequently because of the similarity between sheep and human placentae.

**TABLE 2: foetal development during pregnancy**  
(modified from WebMD<sup>21</sup> and the March of Dimes<sup>16</sup> websites)

Time Period		Foetal Development
1st Trimester	1st Month	Limb buds (arms and legs), heart, lungs, neural tube (spinal cord and brain), placenta, jaw, throat, blood cells form; heart starts beating by day 22
	2nd Month	Ears, arms, legs, eyelids, fingers and toes grow; bone begins to replace cartilage; major organ systems are formed but still developing
	3rd Month	Fingernails, toenails, buds in mouth (future teeth), hair, ears, reproductive organs grow; circulatory and urinary systems begin to function
2nd Trimester	4th Month	Eyelids, eyebrows, eyelashes form; nervous system starts to function
	5th Month	Fingerprints and toe prints formed; muscles develop; period of rapid growth
	6th Month	Eyelids part and eyes open
3rd Trimester	7th Month	Fat deposition; fetus responds to light, sound, and pain
	8th Month	Fat deposition; rapid brain growth; lungs still maturing
	9th Month	Lungs mature, reflexes become coordinated

Foetal sheep whose mothers underwent decompression dives (following U.S. Navy dive tables) sometimes formed bubbles even when the mothers showed no signs of DCI<sup>13,21</sup>.

When the ewes did develop signs of DCI, the foetuses demonstrated even more dramatic evidence of affliction. Researchers reported being able to tell that a foetus had bubbles by detecting early cardiac arrhythmias<sup>21</sup>. For the foetus, these abnormal heartbeats could be life-threatening. The offspring of some sheep that were dived late in pregnancy showed limb weakness and spinal defects associated with DCI, even when the mother had remained symptom-free<sup>21</sup>.

Scientists have long known that so-called 'silent bubbles' - those not associated with symptoms - can develop after diving (note: Dr. Albert Behnke, a pioneer in modern diving medicine and physiology research, is credited for coining this term<sup>4</sup>). Fully functional lungs are extremely effective in filtering bubbles from the circulation. In the foetus, however, most blood bypasses the lungs (via the foramen ovale and ductus arteriosus

shunts), and gas exchange occurs through the placenta. Thus, pulmonary filtration of bubbles does not occur within the foetus. This may increase the risk of arterial gas embolism (AGE), with potentially devastating consequences.

Foetal circulation requires further consideration. During a series of dives that exposed ewes to 100 percent oxygen at 3.0 ATA for approximately 50 minutes, researchers noticed that the circulatory shunts began to close while at depth. Flow through the foramen ovale dropped by 50 percent, and the ductus arteriosus flow fell to zero or even reversed direction<sup>2</sup>.

Once the dives were completed, the circulation reverted to its usual form, and the researchers did not notice any negative effects from the temporary change. Whether the foetus suffered consequences that were not obvious to the researchers was unclear.

The animal study data can be compared with human experience. Premature closure of the ductus arteriosus during human pregnancy has been associated with congestive heart failure<sup>1,3,18</sup> and neonatal death<sup>3,5,18</sup>. Such closure can

unintentionally be induced by prolonged use of indomethacin, a drug commonly used to halt premature labor<sup>9</sup>. Whether scuba diving could induce problematic closure is uncertain, but the possibility should be considered.

### Practical considerations

In addition to possible risk to the foetus, changes in a woman's body during pregnancy might make diving more problematic. Swelling of the mucous membranes in the sinuses could make ear clearing difficult<sup>8,10</sup>, and nausea may increase discomfort<sup>10,27</sup>.

The physical aspects must also be appreciated. A woman's growing abdomen could pose a problem in fitting suits, buoyancy compensation devices, weight belts and other equipment. In addition to the hazards inherent in poorly fitted gear, diving simply may not be enjoyable.

### Decisions

Sifting through the published literature reveals why there is debate over the topic. Data are limited and, in many cases, apparently inconsistent. While this makes drawing conclusions more difficult, it should not be surprising.

Science is very rarely as clear-cut as might be desired. It is difficult to design an ethical experiment that tests only the variable of interest and controls for all others. It is the researcher's job to design the best experiments possible, and it is the individual's or advocate's responsibility to examine the results and decide how to best respond to them.

The question of diving and pregnancy is a difficult one to study since the trend is for women to refrain from diving while pregnant<sup>24</sup>. Most physicians treat diving as they would any drug for which the evidence with respect to pregnancy is incomplete: If there is not a good reason to take it, avoid it.

Anyone who inadvertently dives while pregnant, however, may take solace in the anecdotal evidence from women reporting repeated diving during pregnancy without complication. There is certainly insufficient evidence to warrant termination of a pregnancy. Moreover, if emergency hyperbaric oxygen is required during pregnancy, for example to treat carbon monoxide poisoning, the evidence suggests that the risk to the foetus with treatment is lower than without.

The overall picture of the literature indicates that, while the effect may be small, diving during pregnancy does increase the risk to the foetus, and the consequences could be devastating to all involved. Appreciating these essential factors, the prudent course is to avoid diving while pregnant. While it is possible that some diving could be completed without impact, the absolute risk of any given exposure cannot be determined from the available data. Given the ethical challenges of research on diving during pregnancy and the fact that diving represents a completely avoidable risk for most women, it is unlikely that studies will be conducted to establish the absolute risk in the foreseeable future.

### About the Authors

Heather Held completed an undergraduate degree in biomedical science at Texas A&M University in December 2005. She was placed at the Centre for Hyperbaric Medicine and Environmental Physiology, Duke University Medical Center, Durham, N.C., as a DAN research intern in the summer of 2006.

Dr. Neal Pollock is a research physiologist at the Centre for Hyperbaric Medicine and Environmental Physiology, Duke University Medical Center, Durham, N.C.

### References

1. Arcilla RA, Thilenius OG, Ranniger K. Congestive heart failure from suspected ductal closure in utero. *Journal of Pediatrics* 1969;75(1):74-8.
2. Assali NS, Kirschbaum TH, Dilts PV Jr. Effects of hyperbaric oxygen on uteroplacental and fetal circulation. *Circulation Research* 1968;22(5):573-88.
3. Becker AE, Becker MJ, Wagenvoort CA. Premature contraction of the ductus arteriosus: a cause of foetal death. *Journal of Pathology* 1977;121(2):187-91.
4. Behnke AR. Decompression sickness: advances and interpretations. *Aerospace Medicine* 1971; 42(3):255-267.
5. Ben-David Y, Hallak M, Rotschild A, Sorokin Y, Auslender R, Abramovici H. Indomethacin and fetal ductus arteriosus: complete closure after cessation of prolonged therapeutic course. *Fetal Diagnosis and Therapy* 1996;11(5):341-4.
6. Bolton ME, Alamo AL. Lack of teratogenic effects of air at high ambient pressure in rats. *Teratology* 1981;24:181-5.
7. Bolton ME. Scuba diving and fetal well-being: a survey of 208 women. *Undersea Biomedical Research* 1980;7(3):183-9.
8. Breathnach F, Geoghegan T, Daly S, Turner MJ. Air travel in pregnancy: the air-born study. *Irish Medical Journal* 2004;97(6):167-8.
9. Briggs GG, Freeman RK, Yaffe SJ. *Drugs in Pregnancy and Lactation*. 6th ed. Philadelphia: Lippincott Williams & Wilkins; 2002. p. xxiv, 705/i-13/i.
10. Camporesi EM. Diving and pregnancy. *Seminars in Perinatology* 1996;20(4):292-302.
11. Ferm VH. Teratogenic effects of hyperbaric oxygen. *Proceedings of the society for experimental biology and medicine* 1964;116:975-6.
12. Ferm, VH. The use of the golden hamster in experimental teratology. *Lab Animal Care* 1967;17(5):452-62.
13. Fife WP, Simmang C, Kitzman JV. Susceptibility of fetal sheep to acute decompression sickness. *Undersea Biomedical Research* 1978;5(3):287-92.
14. Fujikura T. Retrorenal fibroplasia and prematurity in newborn rabbits induced by maternal hyperoxia. *American Journal of Obstetrics and Gynecology* 1964;90(7): 854-8.
15. Gilman SC, Bradley ME, Greene KM, Fischer GJ. Fetal development: effects of decompression sickness and treatment. *Aviation, Space, and Environmental Medicine* 1983;54(11):1040-2.
16. Gilman SC, Greene KM, Bradley ME, Biersner RJ. Fetal development: effects of simulated diving and hyperbaric oxygen treatment. *Undersea Biomedical Research* 1982;9(4):297-304.
17. How your baby grows [Online]. 2006 [cited 2006 Jul 11]; Available from: URL: [http://www.marchofdimess.com/pnhec/1808\\_2134.asp](http://www.marchofdimess.com/pnhec/1808_2134.asp)
18. Harlass FE, Duff P, Brady K, Read J. Hydrops fetalis and premature closure of the ductus arteriosus: a review. *Obstetrical and Gynecological Survey* 1989;44(7): 541-3.
19. Joiner JT, editor. *NOAA diving manual*. 4th ed. Flagstaff: Best Publishing Company; 2001.
20. Miller PD, Telford IR, Haas GR. Effect of hyperbaric oxygen on cardiogenesis in the rat. *Biology of the Neonate* 1971;17(1):44-52.
21. Powell MR, Smith MT. Fetal and maternal bubbles detected noninvasively in sheep and goats following hyperbaric decompression. *Undersea Biomedical Research* 1985;12(1):59-67.
22. Shumand T, editor. *Pregnancy: your baby's growth and development* [Online]. 2005 Nov [cited 2006 Jul 11]; Available from: URL: <http://www.webmd.com/content/article/51/40829.htm> <http://www.webmd.com/content/article/51/40805.htm> <http://www.webmd.com/content/article/51/40817.htm>
23. Spencer MP. Decompression limits for compressed air determined by ultrasonically detected blood bubbles. *Journal of Applied Physiology* 1976;40(2):229-35.
24. St Leger Dowse M, Gunby A, Moncad R, Fife C, Bryson P. Scuba diving and pregnancy: can we determine safe limits? *Journal of Obstetrics and Gynaecology* 2006;26(6):509-13.
25. Stock MK, Lanphier EH, Anderson DF, Anderson LC, Phernetton TM, Rankin JHG. Responses of fetal sheep to simulated no-decompression dives. *Journal of Applied Physiology* 1980;40(5):776-80.
26. Telford IR, Miller PD, Haas GF. Hyperbaric oxygen causes fetal wastage in rats. *Lancet* 1969; 2:220-1.
27. Ugucioni DM, Moon RE, Taylor MB. DAN explores fitness and diving issues for women. In: *The DAN guide to dive medical frequently asked questions (FAQs)*. Durham: Divers Alert Network; 2003:209-15.
28. Willson JR, Blessed WB, Blackburn PJ. Hyperbaric exposure during pregnancy in sheep: staged and rapid decompression. *Undersea Biomedical Research* 1983;10(1):11-5.